

Colonoscopy Preparation Using 3 Day Prep



One Week Prior to Colonoscopy

- Stop all **iron supplements** and **vitamins** containing iron 7 days prior to procedure.
- Stop all blood thinning medications as directed by your physician.
- If you take medication for diabetes, adjustments will need to be made in your medication due to changes to your diet. Your doctor will instruct you on medication changes.

Purchase from Pharmacy

- 1 (one) 10 oz. bottle of magnesium citrate
- Milk of magnesia (3 Tbsps.)
- Miralax 238 gram bottle (8.3 oz)
- Dulcolax laxative tablets (4 tablets)
- Gatorade 64 oz. bottle (**NO RED OR PURPLE**)

All items can be bought over the counter. You may purchase plain or aloe wipes to prevent a sore bottom.

Three days prior to procedure Date: _____

- You will be on a **low fiber diet** (no nuts, corn, seeds, or fruit containing seeds). Take 3 tablespoons of Milk of Magnesia at 7 p.m. followed with an 8 oz. glass of water.

Two days prior to procedure Date: _____

- You will be on a **full liquid diet** (Instructions for a full liquid diet are on next page). Drink the 10 oz. bottle of magnesium citrate at 7 p.m. followed with and 8 oz. glass of water.

Day Before Your Colonoscopy Date: _____

- You will need to be on a **clear liquid diet the entire day** (instructions for a clear liquid diet are on the next page).
- At 3 p.m.**, take 2 Dulcolax tablets with a glass of water.
- At 5 p.m.**, add the entire bottle of Miralax with 64 oz. Gatorade. Mix until well dissolved.
- Drink an 8 oz. glass of the Miralax/Gatorade solution every 15 minutes until the solution is gone. *(If you become bloated or nauseated, you may take a break from drinking for 30 minutes to 1 hour to allow it to move downstream. Then continue to drink at a slower rate.)*
- After drinking all of the Miralax/ Gatorade solution, take 2 more Dulcolax tablets.
- If you are diabetic, it is important to keep a check on your blood sugars. You may need to adjust your medications since you will be decreasing the amount of calories you are taking in. We suggest taking half your usual dosage. Please discuss this with your family doctor prior to the procedure.

Your stools should become watery from 30 minutes to 3 hours after starting colon cleansing. It is important to consume the entire bottle of solution to allow the physician a clear view of your colon.

Colonoscopy Preparation Using 3 Day Prep continued

On the Day of Your Colonoscopy

- You **may not have any food or drink after midnight on the night before** your colonoscopy.
- You may brush your teeth.
- You will need a driver after the procedure** because you will be given medication to make you sleepy during the procedure. If your driver needs to leave after dropping you off, please make sure they leave a working number so we may contact them to pick you up after your procedure.

Full Liquid Diet

- Milk, milk products such as chocolate milk, milkshakes, buttermilk, flavored yogurt (without nuts, seeds, or fruit pieces), hot chocolate or cocoa
- Cream of wheat, oatmeal, grits
- Fruit juice and vegetable juice without pulp
- Cream soups, pureed soups, bouillons, and broths
- Pudding, custards, Jell-O, ice cream, and sherbet
- Anything on clear liquid diet list (below)

Clear Liquid Diet

- Ginger ale, orange, sodas, Gatorade, Kool Aid
- Apple juice, white grape, orange juice, lemonade, limeade (no pulp)
- Water, tea, black coffee (no creamer or milk products)
- Broths or bouillon cubes
- Jell-O and popsicles (**NO RED OR PURPLE**)
- Hard candies